

Shape theory via SVD decomposition I

José A. Díaz-García *

Department of Statistics and Computation
 Universidad Autónoma Agraria Antonio Narro
 25350 Buenavista, Saltillo, Coahuila, Mexico
 E-mail: jadiaz@uaaan.mx

Francisco J. Caro-Lopera

Department of Basic Sciences
 Universidad de Medellín
 Carrera 87 No.30-65, of. 5-103
 Medellín, Colombia
 E-mail: fjcaro@udem.edu.co

Abstract

This work finds the non isotropic noncentral elliptical shape distributions via SVD decomposition in the context of zonal polynomials, avoiding the invariant polynomials and the open problems for their computation. The new shape distributions are easily computable and then the inference procedure is based on exact densities instead of the published approximations and asymptotic densities of isotropic models. An application of the technique is illustrated with a classical landmark data in Biology, for this, three models are proposed, the usual Gaussian and two non Gaussian; the best one is chosen by using a modified BIC criterion.

1 Introduction

The multivariate statistical theory of shape has been studied deeply in the last two decades (Goodall and Mardia (1993), Díaz-García *et al.* (1997), Díaz-García *et al.* (2003), Dryden and Mardia (1998) and the references there in, Caro-Lopera *et al.* (2009), among many others. Most of the works are supported by important restrictions (isotropy) for the covariance matrix and models (Normal) in order to obtain known polynomials (zonal polynomials). A sort of approaches are given for shape theory, via QR (Goodall and Mardia (1993)), SVD (Le and Kendall (1993), Goodall (1991), Díaz-García *et al.* (2003)), affine (Goodall and Mardia (1993), Díaz-García *et al.* (2003), Caro-Lopera *et al.* (2009)).

Avoiding the restrictions of isotropy and normality carry some problems, because integration over Euclidean or affine transformations lead to the apparition of invariant polynomials of Davis (1908) which can not computed for large degrees.

This work finds a sequence of transformations which let the construction of shape densities via the singular value decomposition and based on a non restricted non central and non

*Corresponding author

Key words. Shape theory, non-central and non-isotropic shape density, zonal polynomials.
 2000 Mathematical Subject Classification. Primary 62E15; 60E05; secondary 62H99

isotropy elliptical model. The resulting densities avoids the invariant polynomials and they are set in terms of series of zonal polynomials which can be computed by suitable modifications of the existing algorithms for hypergeometric series (Koev and Edelman (2006)).

The work is structured as follows, the main principle and the size and shape distribution is given in section 2, then the shape density is obtained in section 3; the associated excluding reflection densities are considered in section 4 and finally some particular models are derived as corollaries in section 6 which also presents an application in mouse vertebra by studying three models (the usual Gaussian and two non Gaussian) with the modified BIC criterion.

2 Main principle and SVD size-and-shape density

It is known that the shape of an object is all geometrical information that remains after filtering out translation, rotation and scale information of an original figure (represented by a matrix $\mathbf{X}$) comprised in N landmarks in K dimensions. So, we say that two figures, $\mathbf{X}_1 : N \times K$ and $\mathbf{X}_2 : N \times K$ have the same shape if they are related by a special similarity transformation $\mathbf{X}_2 = \beta \mathbf{X}_1 \mathbf{H} + \mathbf{1}_N \gamma'$, where $\mathbf{H} : K \times K \in SO(K)$ (the rotation), $\gamma : K \times 1$ (the translation), $\mathbf{1}_N : N \times 1$, $\mathbf{1}_N = (1, 1, \dots, 1)'$, and $\beta > 0$ (the scale). Thus, in this context, the shape of a matrix $\mathbf{X}$ is all the geometrical information about $\mathbf{X}$ that is invariant under Euclidean similarity transformations.

Now, multivariate statistical theory of shape compares shapes of objects in presence of randomness, so if we assume that a figure $\mathbf{X}$, comprised in N landmarks in K , follows an elliptical distribution $\mathbf{X} \sim \mathcal{E}_{N \times K}(\mu_{\mathbf{x}}, \Sigma_{\mathbf{x}}, \Theta, h)$, it is of interest to remove translation, scaling, rotation from $\mathbf{X}$. Clearly, the sequence $\mathbf{LX} = \mathbf{Y} = \mathbf{H}'\mathbf{D}\mathbf{P} = r\mathbf{W}(\mathbf{u})\mathbf{P}$ removes the translation (by a sub Helmert matrix $\mathbf{L}$, for example), the rotation (by the SVD of $\mathbf{Y}$) and the scale (by dividing for the norm of $\mathbf{Y}$). In order to obtain the density of $\mathbf{W}$ we need to integrate over the similarity group; it is easy to see that the elliptical assumption lead to the product of two traces which irremediably expands in terms of invariant polynomials of two matrix arguments (Davis (1908)), and the shape densities are not computable for large degrees.

So the classical statistical multivariate analysis restricts the models for the original landmark data in order to obtain densities which are expanded in terms of studied polynomials such as the zonal polynomials which are computable (Goodall and Mardia (1993)), otherwise, as we proved in the last sentence, the densities involve non computable polynomials for large degrees.

From the practical point of view the restrictions affect the applications; i.e., the isotropic assumption $\Theta = \mathbf{I}_K$ for an elliptical shape model of the form

$$\mathbf{X} \sim \mathcal{E}_{N \times K}(\mu_{\mathbf{x}}, \Sigma_{\mathbf{x}}, \Theta, h),$$

restricts substantially the correlations of the landmarks in the figure, specially in objects with symmetries as in the case of mouse vertebra for example, among many others (Dryden and Mardia (1998)). So, we expect the non isotropic model, with any positive definite matrix Θ , as the best model for considering all the possible correlations among the anatomical (geometrical or mathematical) points. However, using the classical approach of the published literature of shape (see for example Goodall and Mardia (1993)) under the non isotropic model, we obtain immediately invariant polynomials, which can not be computed at this time for large degrees.

In order to avoid this problem consider the following procedure: Let

$$\mathbf{X} \sim \mathcal{E}_{N \times K}(\mu_{\mathbf{x}}, \Sigma_{\mathbf{x}}, \Theta, h),$$

if $\Theta^{1/2}$ is the positive definite square root of the matrix Θ , i.e. $\Theta = (\Theta^{1/2})^2$, with $\Theta^{1/2} : K \times K$, Gupta and Varga (1993, p. 11), and noting that

$$\mathbf{X}\Theta^{-1}\mathbf{X}' = \mathbf{X}(\Theta^{-1/2}\Theta^{-1/2})^{-1}\mathbf{X}' = \mathbf{X}\Theta^{-1/2}(\mathbf{X}\Theta^{-1/2})' = \mathbf{Z}\mathbf{Z}',$$

where

$$\mathbf{Z} = \mathbf{X}\Theta^{-1/2},$$

then

$$\mathbf{Z} \sim \mathcal{E}_{N \times K}(\boldsymbol{\mu}_{\mathbf{Z}}, \boldsymbol{\Sigma}_{\mathbf{Z}}, \mathbf{I}_K, h),$$

with $\boldsymbol{\mu}_{\mathbf{Z}} = \boldsymbol{\mu}_{\mathbf{X}}\Theta^{-1/2}$, (see Gupta and Varga (1993, p. 20)).

And we arrive at the classical starting point in shape theory where the original landmark matrix is replaced by $\mathbf{Z} = \mathbf{X}\Theta^{-1/2}$. Then we can proceed as usual, removing from $\mathbf{Z}$, translation, scale, rotation and/or reflection in order to obtain the shape of $\mathbf{Z}$ (or $\mathbf{X}$) via the SVD decomposition, for example.

The SVD decomposition has two version in shape theory, Goodall (1991) and Le and Kendall (1993), we focus in this paper on Goodall's approach.

Let $n = \min(m, K)$, $\mathbf{Y} = \mathbf{H}'\mathbf{D}\mathbf{P}$ be the nonsingular part of the SVD, where $\mathbf{H} : n \times m$, $\mathbf{H} \in \mathcal{V}_{n,m}$ represents the Stiefel manifold, $\mathbf{D} = \text{diag}(D_1, D_2, \dots, D_n)$ with $D_1 \geq D_2 \geq \dots \geq D_n$ and $\mathbf{P} : n \times K$, $\mathbf{P} \in \mathcal{V}_{n,K}$.

Thus the SVD shape coordinates $\mathbf{u}$ of $\mathbf{X}$ may be found by the following procedure

$$\mathbf{L}\mathbf{X}\Theta^{-1/2} = \mathbf{L}\mathbf{Z} = \mathbf{Y} = \mathbf{H}'\mathbf{D}\mathbf{P} = r\mathbf{W}\mathbf{P} = r\mathbf{W}(\mathbf{u})\mathbf{P} \quad (1)$$

where the SVD shape coordinate system is given by $\mathbf{H}'\mathbf{D}$ (Goodall (1991, pp. 296-298)) and $r = \|\mathbf{H}\mathbf{D}\| = (\text{tr } \mathbf{D}\mathbf{H}\mathbf{H}'\mathbf{D})^{1/2} = (\text{tr } \mathbf{D}^2)^{1/2} = \|\mathbf{D}\| = \|\mathbf{Y}\|$. Before defining $\mathbf{W}$ and $\mathbf{u}$, note that when $n = K$ two cases may be distinguished.

1. $\mathbf{P}$ includes reflection, $\mathbf{P} \in \mathcal{O}(K)$, $|\mathbf{P}| = \pm 1$, $\mathbf{D}_K \geq 0$ and $(\mathbf{H}, \mathbf{D})$, written $(\mathbf{H}, \mathbf{D})^R$ for definiteness, contains reflection SVD shape co-ordinates.
2. $\mathbf{P}$ excludes reflection, $\mathbf{P} \in \mathcal{SO}(K)$, $|\mathbf{P}| = +1$, $|D_K| \geq 0$, $\text{sign}(D_m) = \text{sign}|\mathbf{X}|$ and $(\mathbf{H}, \mathbf{D})$, may be written $(\mathbf{H}, \mathbf{D})^{NR}$ for definiteness.

Now the SVD shape matrix $\mathbf{W}$ is obtained by dividing the $\mathbf{H}'\mathbf{D}$ matrix by r , when $\mathbf{W}$ may include or exclude reflection, in which case we obtain, respectively, $\mathbf{W}^R = (\mathbf{H}'\mathbf{D})^R/r$ or $\mathbf{W}^{NR} = (\mathbf{H}'\mathbf{D})^{NR}/r$. Finally $\mathbf{u}$ is composed of the $mn - 1$ generalized polar coordinates.

Our interest now lies in finding the corresponding densities associated with the process described in (1). Thus we obtain the joint density of $(\mathbf{H}, \mathbf{D})$ and the density of $\mathbf{W}(\mathbf{u})$.

In order to obtain the size and shape density we need some integrals involving zonal polynomials, extending James (1964, eq. (22)).

Lemma 2.1. *Let $\mathbf{X} : K \times n$, $\mathbf{Y} : K \times K$ and $\mathbf{H} \in \mathcal{V}_{n,K}$. Then*

1.

$$\int_{\mathbf{H} \in \mathcal{V}_{n,K}} [\text{tr}(\mathbf{Y} + \mathbf{X}\mathbf{H})]^p (\mathbf{H}\mathbf{d}\mathbf{H}') = \frac{2^n \pi^{Kn/2}}{\Gamma_n[\frac{1}{2}K]} \sum_{f=0}^{\infty} \sum_{\lambda} \frac{(p)_{2f} (\text{tr } \mathbf{Y})^{p-2f}}{(\frac{1}{2}K)_{\lambda}} \frac{C_{\lambda}(\frac{1}{4}\mathbf{X}\mathbf{X}')}{f!}$$

where $|(\text{tr } \mathbf{Y})^{-1} \text{tr } \mathbf{X}\mathbf{H}| < 1$ and $\text{tr } \mathbf{Y} \neq 0$.

2. $\int_{\mathbf{H} \in \mathcal{V}_{n,K}} \text{tr}(\mathbf{Y} + \mathbf{X}\mathbf{H}) \text{etr}\{r(\mathbf{Y} + \mathbf{X}\mathbf{H})\} (\mathbf{H}\mathbf{d}\mathbf{H}') =$

$$\frac{2^n \pi^{Kn/2}}{\Gamma_n[\frac{1}{2}K]} \text{etr}\{r\mathbf{Y}\} \left\{ \text{tr } \mathbf{Y}_0 F_1\left(\frac{1}{2}K; \frac{r^2}{4}\mathbf{X}\mathbf{X}'\right) + \sum_{f=0}^{\infty} \sum_{\lambda} \frac{(f + \frac{1}{2})}{(\frac{1}{2}K)_{\lambda}} \frac{C_{\lambda}(\frac{1}{4}\mathbf{X}\mathbf{X}')}{f!} \right\},$$

where $p \in \mathbb{R}$, $r \in \mathbb{R}$, $C_\kappa(\mathbf{B})$ are the zonal polynomials of $\mathbf{B}$ corresponding to the partition $\kappa = (f_1, \dots, f_p)$ of f , with $\sum_{i=1}^p f_i = f$; and $(a)_\kappa = \prod_{i=1}^p (a - (j-1)/2)_{f_j}$, $(a)_f = a(a+1) \cdots (a+f-1)$, are the generalized hypergeometric coefficients and ${}_0F_1$ is the Bessel function, James (1964).

Proof.

1. From Lemma 9.5.3 Muirhead (1982, Lemma 9.5.3, p. 397) we have

$$\int_{\mathbf{H} \in \mathcal{V}_{n,K}} [\text{tr}(\mathbf{Y} + \mathbf{X}\mathbf{H})]^p (\mathbf{H} d\mathbf{H}') = \frac{2^n \pi^{Kn/2}}{\Gamma_n[\frac{1}{2}K]} \int_{\mathcal{O}(K)} [\text{tr}(\mathbf{Y} + \mathbf{X}\mathbf{H})]^p (d\mathbf{H}).$$

Furthermore, for $\text{tr } \mathbf{Y} \neq 0$ and $|(\text{tr } \mathbf{Y})^{-1} \text{tr } \mathbf{X}\mathbf{H}| < 1$

$$[\text{tr}(\mathbf{Y} + \mathbf{X}\mathbf{H})]^p = (\text{tr } \mathbf{Y})^p \sum_{f=0}^{\infty} \frac{(p)_f}{f!} (\text{tr } \mathbf{Y})^{-f} (\text{tr } \mathbf{X}\mathbf{H})^f.$$

Now from James (1964, eqs. (46) and (22)) it follows that

$$\begin{aligned} \int_{\mathbf{H} \in \mathcal{V}_{n,K}} [\text{tr}(\mathbf{Y} + \mathbf{X}\mathbf{H})]^p (\mathbf{H} d\mathbf{H}') &= \\ \frac{2^n \pi^{Kn/2}}{\Gamma_n[\frac{1}{2}K]} \sum_{f=0}^{\infty} \sum_{\lambda} \frac{(p)_{2f} (\text{tr } \mathbf{Y})^{-2f}}{(2f)!} \frac{(\frac{1}{2})_f}{(\frac{1}{2}K)_\lambda} C_\lambda(\mathbf{X}\mathbf{X}'), \end{aligned}$$

the result follows, noting that $(\frac{1}{2})_f / (2f)! = 1/(4^f f!)$ and that $C_\lambda(a\mathbf{X}\mathbf{X}') = a^f C_\lambda(\mathbf{X}\mathbf{X}')$.

2. This follows by expanding the exponentials in series of powers and by applying (22) and (27) from James (1964). $\square$

Now, the jacobian of the corresponding decomposition is provided next:

Lemma 2.2. *Let be $\mathbf{Y} : N-1 \times K$, then there exist $\mathbf{V} \in \mathcal{V}_{n,N-1}$, $\mathbf{H} \in \mathcal{V}_{n,K}$ and $\mathbf{D} : n \times n$, $\mathbf{D} = \text{diag}(D_1, \dots, D_n)$, $n = \min(N-1, K)$; $D_1 \geq D_2 \geq \dots \geq D_n \geq 0$, such that $\mathbf{Y} = \mathbf{V}'\mathbf{D}\mathbf{H}$; This factorization is termed non-singular part of the SVD. Then*

$$(d\mathbf{Y}) = 2^{-n} |\mathbf{D}|^{N-1+K-2n} \prod_{i < j}^n (D_i^2 - D_j^2) (d\mathbf{D}) (\mathbf{V} d\mathbf{V}') (\mathbf{H} d\mathbf{H}').$$

Proof. See Díaz-García *et al.* (1997). $\square$

So, we can obtain:

Theorem 2.1. *The joint density of $(\mathbf{V}, \mathbf{D})$ is*

$$\begin{aligned} f_{\mathbf{V}, \mathbf{D}}(\mathbf{V}, \mathbf{D}) &= \frac{\pi^{\frac{nK}{2}} |\mathbf{D}|^{N-1+K-2n} \prod_{i < j} (D_i^2 - D_j^2)}{\Gamma_n[\frac{K}{2}] |\Sigma|^{\frac{K}{2}}} \\ &\times \sum_{t=0}^{\infty} \sum_{\kappa} \frac{h^{(2t)} [\text{tr}(\Sigma^{-1} \mathbf{V}' \mathbf{D}^2 \mathbf{V} + \Omega)]}{t! (\frac{1}{2}K)_\kappa} C_\kappa(\Omega \Sigma^{-1} \mathbf{V}' \mathbf{D}^2 \mathbf{V}). \end{aligned}$$

Proof. Let be $\mathbf{\Omega} = \mathbf{\Sigma}^{-1} \boldsymbol{\mu} \boldsymbol{\Theta}^{-1} \boldsymbol{\mu}'$, so the density of $\mathbf{Y}$ is given by

$$f_{\mathbf{Y}}(\mathbf{Y}) = \frac{1}{|\mathbf{\Sigma}|^{\frac{K}{2}}} h \left[\text{tr} (\mathbf{\Sigma}^{-1} \mathbf{Y} \mathbf{Y}' + \mathbf{\Omega}) - 2 \text{tr} \boldsymbol{\mu}' \mathbf{\Sigma}^{-1} \mathbf{Y} \right].$$

Now, make the change of variables $\mathbf{Y} = \mathbf{V}' \mathbf{D} \mathbf{H}$, so, by Lemma 2.2, the joint density function of $\mathbf{V}$, $\mathbf{D}$, $\mathbf{H}$ is

$$\begin{aligned} dF_{\mathbf{V}, \mathbf{D}, \mathbf{H}}(\mathbf{V}, \mathbf{D}, \mathbf{H}) &= \frac{2^{-n} |\mathbf{D}|^{N-1+K-2n} \prod_{i < j} (D_i^2 - D_j^2)}{|\mathbf{\Sigma}|^{\frac{K}{2}}} (\mathbf{V} d\mathbf{V}') (d\mathbf{D}) \\ &\times h \left[\text{tr} (\mathbf{\Sigma}^{-1} \mathbf{V}' \mathbf{D}^2 \mathbf{V} + \mathbf{\Omega}) - 2 \text{tr} \boldsymbol{\mu}' \mathbf{\Sigma}^{-1} \mathbf{V}' \mathbf{D} \mathbf{H} \right] (\mathbf{H} d\mathbf{H}'). \end{aligned}$$

Expanding in power series

$$\begin{aligned} dF_{\mathbf{V}, \mathbf{D}, \mathbf{H}}(\mathbf{V}, \mathbf{D}, \mathbf{H}) &= \frac{2^{-n} |\mathbf{D}|^{N-1+K-2n} \prod_{i < j} (D_i^2 - D_j^2)}{|\mathbf{\Sigma}|^{\frac{K}{2}}} (\mathbf{V} d\mathbf{V}') (d\mathbf{D}) \\ &\times \sum_{t=0}^{\infty} \frac{1}{t!} h^{(t)} \left[\text{tr} (\mathbf{\Sigma}^{-1} \mathbf{V}' \mathbf{D}^2 \mathbf{V} + \mathbf{\Omega}) \right] \left[\text{tr} (-2 \text{tr} \boldsymbol{\mu}' \mathbf{\Sigma}^{-1} \mathbf{V}' \mathbf{D} \mathbf{H}) \right]^t (\mathbf{H} d\mathbf{H}'). \end{aligned}$$

From Lemma 2.1

$$\int_{\mathcal{V}_{n,K}} \left[\text{tr} (-2 \text{tr} \boldsymbol{\mu}' \mathbf{\Sigma}^{-1} \mathbf{V}' \mathbf{D} \mathbf{H}) \right]^{2t} (\mathbf{H} d\mathbf{H}') = \frac{2^n \pi^{\frac{nK}{2}}}{\Gamma_n \left[\frac{K}{2} \right]} \sum_{\kappa} \frac{\left(\frac{1}{2} \right)_t 4^t}{\left(\frac{1}{2} K \right)_{\kappa}} C_{\kappa} (\mathbf{\Omega} \mathbf{\Sigma}^{-1} \mathbf{V}' \mathbf{D}^2 \mathbf{V}).$$

Observing that $\frac{\left(\frac{1}{2} \right)_t 4^t}{(2t)!} = \frac{1}{t!}$, the marginal joint density of $\mathbf{V}$, $\mathbf{D}$ is given by

$$\begin{aligned} dF_{\mathbf{V}, \mathbf{D}}(\mathbf{V}, \mathbf{D}) &= \frac{\pi^{\frac{nK}{2}} |\mathbf{D}|^{N-1+K-2n} \prod_{i < j} (D_i^2 - D_j^2)}{\Gamma_n \left(\frac{K}{2} \right) |\mathbf{\Sigma}|^{\frac{K}{2}}} \\ &\times \sum_{t=0}^{\infty} \frac{h^{(2t)} \left[\text{tr} (\mathbf{\Sigma}^{-1} \mathbf{V}' \mathbf{D}^2 \mathbf{V} + \mathbf{\Omega}) \right]}{t! \left(\frac{1}{2} K \right)_{\kappa}} C_{\kappa} (\mathbf{\Omega} \mathbf{\Sigma}^{-1} \mathbf{V}' \mathbf{D}^2 \mathbf{V}) (\mathbf{V} d\mathbf{V}') (d\mathbf{D}). \quad \square \end{aligned}$$

Now, let be

$$\mathbf{R}' = \mathbf{V}' \mathbf{D} \tag{2}$$

then $\mathbf{V}' = \mathbf{R}' \mathbf{D}^{-1}$, so $d\mathbf{V} = d\mathbf{R}' \mathbf{D}^{-1}$ and $\mathbf{V} d\mathbf{V}' = \mathbf{D}^{-1} \mathbf{R} d\mathbf{R}' \mathbf{D}^{-1}$. But $\mathbf{R} d\mathbf{R}' : n \times n$ is skew symmetric, thus

$$(\mathbf{V} d\mathbf{V}') = |\mathbf{D}|^{-n+1} (\mathbf{R} d\mathbf{R}') \tag{3}$$

Theorem 2.2. *The SVD reflection size-and-shape density is*

$$\begin{aligned} dF_{\mathbf{R}}(\mathbf{R}) &= \frac{2^{-n} \Gamma_n \left[\frac{N+K-2n}{2} \right] \pi^{\frac{nK}{2}}}{\pi^{n(N+K-(3n-1)/2)} \Gamma_n \left[\frac{K}{2} \right] |\mathbf{\Sigma}|^{\frac{K}{2}}} (d\mathbf{D}) \\ &\times \sum_{t=0}^{\infty} \frac{h^{(2t)} \left[\text{tr} (\mathbf{\Sigma}^{-1} \mathbf{R}' \mathbf{R} + \mathbf{\Omega}) \right]}{t! \left(\frac{1}{2} K \right)_{\kappa}} C_{\kappa} (\mathbf{\Omega} \mathbf{\Sigma}^{-1} \mathbf{R}' \mathbf{R}) (\mathbf{R} d\mathbf{R}'). \end{aligned} \tag{4}$$

Proof. The joint density function of $\mathbf{V}, \mathbf{D}$ is

$$\begin{aligned} dF_{\mathbf{V}, \mathbf{D}}(\mathbf{V}, \mathbf{D}) &= \frac{\pi^{\frac{nK}{2}} |\mathbf{D}|^{N-1+K-2n} \prod_{i < j} (D_i^2 - D_j^2)}{\Gamma_n \left[\frac{K}{2} \right] |\Sigma|^{\frac{K}{2}}} (d\mathbf{D}) \\ &\times \sum_{t=0}^{\infty} \sum_{\kappa} \frac{h^{(2t)} [\text{tr} (\Sigma^{-1} \mathbf{V}' \mathbf{D}^2 \mathbf{V} + \Omega)] C_{\kappa} (\Omega \Sigma^{-1} \mathbf{V}' \mathbf{D} \mathbf{V})}{t! \left(\frac{1}{2} K \right)_{\kappa}} (\mathbf{V} d\mathbf{V}'). \end{aligned}$$

Now, let be $\mathbf{R}' = \mathbf{V}' \mathbf{D}$ then $\mathbf{V}' = \mathbf{R}' \mathbf{D}^{-1}$, so $d\mathbf{V}' = d\mathbf{R}' \mathbf{D}^{-1}$ and $\mathbf{V} d\mathbf{V}' = \mathbf{D}^{-1} \mathbf{R} d\mathbf{R}' \mathbf{D}^{-1}$, where $\mathbf{R} d\mathbf{R}'$ is an $n \times n$ skew-symmetric matrix.

$$(\mathbf{V} d\mathbf{V}') = |\mathbf{D}^{-1}|^{n-1} (\mathbf{R} d\mathbf{R}') = |\mathbf{D}|^{-n+1} (\mathbf{R} d\mathbf{R}').$$

Thus the joint density function of $\mathbf{R}, \mathbf{D}$ is

$$\begin{aligned} dF_{\mathbf{R}, \mathbf{D}}(\mathbf{R}, \mathbf{D}) &= \frac{\pi^{\frac{nK}{2}} |\mathbf{D}|^{N+K-3n} \prod_{i < j} (D_i^2 - D_j^2)}{\Gamma_n \left[\frac{K}{2} \right] |\Sigma|^{\frac{K}{2}}} (d\mathbf{D}) \\ &\times \sum_{t=0}^{\infty} \sum_{\kappa} \frac{h^{(2t)} [\text{tr} (\Sigma^{-1} \mathbf{R}' \mathbf{R} + \Omega)] C_{\kappa} (\Omega \Sigma^{-1} \mathbf{R}' \mathbf{R})}{t! \left(\frac{1}{2} K \right)_{\kappa}} (\mathbf{R} d\mathbf{R}'). \end{aligned}$$

For the integration with respect to $\mathbf{D}$ note that if $\mathbf{D}^2 = \mathbf{L}$, so $dL_i = 2D_i dD_i$ and $(d\mathbf{L}) = 2^n |\mathbf{D}| (d\mathbf{D})$, thus

$$(d\mathbf{D}) = 2^{-n} |\mathbf{L}|^{-\frac{1}{2}} (d\mathbf{L}).$$

Let be

$$J = \int_{\mathbf{D}} |\mathbf{D}|^{N+K-3n} \prod_{i < j} (D_i^2 - D_j^2) (d\mathbf{D})$$

so

$$\begin{aligned} J &= \int_{\mathbf{L}} |\mathbf{L}^{\frac{1}{2}}|^{N+K-3n} \prod_{i < j} (L_i - L_j) 2^{-n} |\mathbf{L}|^{-\frac{1}{2}} (d\mathbf{L}) \\ &= 2^{-n} \int_{\mathbf{L}} |\mathbf{L}|^{\frac{1}{2}(N+K-3n-1)} \prod_{i < j} (L_i - L_j) (d\mathbf{L}) \end{aligned}$$

From Fang and Zhang (1990), eq. (3.29), p.102. we have that

$$\begin{aligned} J &= \frac{\prod_{i=1}^n \Gamma \left[\frac{1}{2} (N + K - 2n - i + 1) \right]}{\pi^{\frac{1}{2}n(N+K-2n) + \frac{1}{2}n}} \frac{\pi^{\frac{1}{2}n(N+K-2n) + \frac{1}{2}}}{\prod_{i=1}^n \Gamma \left[\frac{1}{2} (N + K - 2n - i + 1) \right]} \\ &\times \int_{\mathbf{L}} |\mathbf{L}|^{[(N+K-2n)-n-1]/2} \prod_{i < j} (L_i - L_j) (d\mathbf{L}) \\ &= \frac{2^{-n} \pi^{n(n-1)/4} \prod_{i=1}^n \Gamma \left[\frac{1}{2} (N + K - 2n - i + 1) \right]}{\pi^{\frac{n}{2}(N+K-2n) + \frac{n(n-1)}{4}}} \\ &= \frac{\Gamma_n \left[\frac{1}{2} (N + K - 2n) \right] 2^{-n}}{\pi^{\frac{n}{2}(N+K-2n) - \frac{3n-1}{2}}}, \end{aligned}$$

then

$$dF_{\mathbf{R}}(\mathbf{R}) = \frac{\Gamma_n \frac{N+K-2n}{2} 2^{-n} \pi^{\frac{nK}{2}}}{\Gamma_n \left[\frac{K}{2} \right] \pi^{\frac{n}{2} (N+K-\frac{3n-1}{2})} |\Sigma|^{\frac{K}{2}}} \times \sum_{t=0}^{\infty} \sum_{\kappa} \frac{h^{(2t)} [\text{tr} (\Sigma^{-1} \mathbf{R}' \mathbf{R} + \Omega)] C_{\kappa} (\Omega \Sigma^{-1} \mathbf{R}' \mathbf{R})}{t! \left(\frac{1}{2} K \right)_{\kappa}} (\mathbf{R} d\mathbf{R}') \quad \square$$

3 Reflection Shape Density

For the SVD reflection shape density consider the following transformations

$$\mathbf{L} \mathbf{X} \Theta^{-1/2} = \mathbf{L} \mathbf{Z} = \mathbf{Y} = (\mathbf{V}' \mathbf{D}) \mathbf{H} \equiv \mathbf{R}' \mathbf{H} = r \mathbf{W} \mathbf{H} = r \mathbf{W}(\mathbf{u}) \mathbf{H},$$

where $\mathbf{R}' = \mathbf{V}' \mathbf{D}$ and $\mathbf{W} = \mathbf{R}'/r$.

Now, note that $\mathbf{V}' \mathbf{D}$ contains $(N-1)n$ coordinates. Then

$$\text{vec } \mathbf{W} = \frac{1}{r} \text{vec}(\mathbf{V}' \mathbf{D}), \quad r = \|\mathbf{V}' \mathbf{D}\| = \sqrt{\text{tr } \mathbf{V}' \mathbf{D}^2 \mathbf{V}} = \|\mathbf{Y}\|.$$

Then, by Muirhead (1982, Theorem 2.1.3, p. 55):

$$\begin{aligned} (d \text{vec } \mathbf{W}(\mathbf{u})) &= r^m \prod_{i=1}^m \sin^{m-i} \theta_i \bigwedge_{i=1}^m d\theta_i \wedge dr \\ &= r^m J(\mathbf{u}) \bigwedge_{i=1}^m d\theta_i \wedge dr, \end{aligned}$$

with $m = (N-1)n - 1$, $\mathbf{u} = (\theta_1, \dots, \theta_m)'$.

Hence,

Theorem 3.1. *The SVD reflection shape density is given by*

$$\begin{aligned} dF_{\mathbf{W}}(\mathbf{W}) &= \frac{\Gamma_n \left[\frac{N+K-2n}{2} \right] J(\mathbf{u}) \pi^{\frac{nK}{2}}}{2^n \pi^{\frac{n}{2} (N+K-\frac{3n-1}{2})} \Gamma_n \left[\frac{K}{2} \right] |\Sigma|^{\frac{K}{2}}} \sum_{t=0}^{\infty} \sum_{\kappa} \frac{C_{\kappa} (\Omega \Sigma^{-1} \mathbf{W}' \mathbf{W})}{t! \left(\frac{1}{2} K \right)_{\kappa}} (\mathbf{W} d\mathbf{W}') \\ &\times \int_0^{\infty} r^{m+n+2t-1} h^{(2t)} [r^2 \text{tr } \Sigma^{-1} \mathbf{W}' \mathbf{W} + \text{tr } \Omega] (dr). \end{aligned} \quad (5)$$

Proof. The density of $\mathbf{R}$ is

$$\begin{aligned} dF_{\mathbf{R}}(\mathbf{R}) &= \frac{\Gamma_n \left[\frac{N+K-2n}{2} \right] 2^{-n} \pi^{\frac{nK}{2}}}{\pi^{\frac{n}{2} [N+K-\frac{3n-1}{2}]} \Gamma_n \left[\frac{K}{2} \right] |\Sigma|^{\frac{K}{2}}} \\ &\times \sum_{t=0}^{\infty} \sum_{\kappa} \frac{h^{(2t)} [\text{tr} (\Sigma^{-1} \mathbf{R}' \mathbf{R} + \Omega)] C_{\kappa} (\Omega \Sigma^{-1} \mathbf{R}' \mathbf{R})}{t! \left(\frac{1}{2} K \right)_{\kappa}} (\mathbf{R} d\mathbf{R}'). \end{aligned}$$

Putting $\mathbf{W}(\mathbf{u}) = \mathbf{R}'/r$, the joint density of r and $\mathbf{u}$ is

$$\begin{aligned} dF_{r, \mathbf{W}(\mathbf{u})}(r, \mathbf{W}(\mathbf{u})) &= \frac{\Gamma_n \left[\frac{N+K-2n}{2} \right] 2^{-n} \pi^{\frac{nK}{2}} r^m J(\mathbf{u})}{\pi^{\frac{n}{2} [N+K-\frac{3n-1}{2}]} \Gamma_n \left[\frac{K}{2} \right] |\Sigma|^{\frac{K}{2}}} (r^2 \mathbf{W} d\mathbf{W}') \\ &\times \sum_{t=0}^{\infty} \sum_{\kappa} \frac{h^{(2t)} [\text{tr} (r^2 \Sigma^{-1} \mathbf{W}' \mathbf{W} + \Omega)] C_{\kappa} (r^2 \Omega \Sigma^{-1} \mathbf{W}' \mathbf{W})}{t! \left(\frac{1}{2} K \right)_{\kappa}}. \end{aligned}$$

Note that

1. $C_\kappa(r^2 \mathbf{\Omega} \mathbf{\Sigma}^{-1} \mathbf{W}' \mathbf{W}) = r^{2t} C_\kappa(\mathbf{\Omega} \mathbf{\Sigma}^{-1} \mathbf{W}' \mathbf{W})$.
2. $(r^2 \mathbf{W} d\mathbf{W}') = ((r\mathbf{I}) \mathbf{W} d\mathbf{W}' (r\mathbf{I})) = |r\mathbf{I}|^{n-1} (\mathbf{W} d\mathbf{W}') = r^{n-1} (\mathbf{W} d\mathbf{W}')$.

Collecting powers of r , the marginal of $\mathbf{W}$ is

$$dF_{\mathbf{W}}(\mathbf{W}) = \frac{\Gamma_n \left[\frac{N+K-2n}{2} \right] 2^{-n} \pi^{\frac{nK}{2}} J(\mathbf{u})}{\pi^{\frac{n}{2} [N+K-\frac{3n-1}{2}]} \Gamma_n \left[\frac{K}{2} \right] |\mathbf{\Sigma}|^{\frac{K}{2}}} (\mathbf{W} d\mathbf{W}') \sum_{t=0}^{\infty} \sum_{\kappa} \frac{C_\kappa(\mathbf{\Omega} \mathbf{\Sigma}^{-1} \mathbf{W}' \mathbf{W})}{t! \left(\frac{1}{2} K \right)_\kappa} \\ \times \int_0^\infty r^{m+n+2t-1} h^{(2t)} [r^2 \text{tr} \mathbf{\Sigma}^{-1} \mathbf{W}' \mathbf{W} + \text{tr} \mathbf{\Omega}] (dr). \quad \square$$

4 Distributions excluding reflection

Recall that the SVD shape coordinates $\mathbf{u}$ of $\mathbf{X}$ are obtained as follows

$$\mathbf{L} \mathbf{X} \mathbf{\Theta}^{-1/2} = \mathbf{L} \mathbf{Z} = \mathbf{Y} = \mathbf{H}' \mathbf{D} \mathbf{P} = r \mathbf{W} \mathbf{P} = r \mathbf{W}(\mathbf{u}) \mathbf{P} \quad (6)$$

where the SVD shape coordinate system is given by $\mathbf{H}' \mathbf{D}$ (Goodall (1991, pp. 296-298)) and $r = \|\mathbf{H} \mathbf{D}\| = (\text{tr} \mathbf{D} \mathbf{H} \mathbf{H}' \mathbf{D})^{1/2} = (\text{tr} \mathbf{D}^2)^{1/2} = \|\mathbf{D}\| = \|\mathbf{Y}\|$. When $n = K$ we studied the distributions including reflection, i.e. $\mathbf{P} \in \mathcal{O}(k)$, $|\mathbf{P}| = \pm 1$, $\mathbf{D}_K \geq 0$ and $(\mathbf{H}, \mathbf{D})$, written $(\mathbf{H}, \mathbf{D})^R$ for definiteness, contains reflection SVD shape co-ordinates.

In this section we consider the case when $\mathbf{P}$ excludes reflection, thus $\mathbf{P} \in \mathcal{SO}(K)$, $|\mathbf{P}| = +1$, $|D_K| \geq 0$, $\text{sign}(D_m) = \text{sign}|\mathbf{X}|$ and $(\mathbf{H}, \mathbf{D})$, may be written $(\mathbf{H}, \mathbf{D})^{NR}$ for definiteness.

Finally, we have that the excluding reflection SVD size-and-shape and SVD shape densities are given by (4) and (5) divided by 2, respectively.

5 Central Case

Now, we can derive easily the corresponding central distributions of this work.

Corollary 5.1. *The central reflection SVD size-and-shape density is*

$$dF_{\mathbf{R}}(\mathbf{R}) = \frac{2^{-n} \Gamma_n \left[\frac{N+K-2n}{2} \right] \pi^{\frac{nK}{2}}}{\pi^{\frac{n}{2} [N+K-\frac{3n-1}{2}]} \Gamma_n \left[\frac{K}{2} \right] |\mathbf{\Sigma}|^{\frac{K}{2}}} h [\text{tr} \mathbf{\Sigma}^{-1} \mathbf{R}' \mathbf{R}] (\mathbf{R} d\mathbf{R}').$$

Proof. Just take $\boldsymbol{\mu} = 0$ in Theorem 2.2 and recall that $h^{(0)}(\cdot) \equiv h(\cdot)$. $\square$

Finally,

Corollary 5.2. *The central reflection SVD shape density is invariant under the elliptical family and it is given by*

$$dF_{\mathbf{W}}(\mathbf{W}) = \frac{2^{-n-1} \Gamma_n \left[\frac{N+K-2n}{2} \right] \pi^{\frac{nK}{2}} \Gamma \left[\frac{m+n}{2} \right]}{\pi^{\frac{n}{2} [N+K-\frac{3n-1}{2}] + \frac{m+n}{2}} \Gamma_n \left[\frac{K}{2} \right] |\mathbf{\Sigma}|^{\frac{K}{2}}} h [\text{tr} \mathbf{\Sigma}^{-1} \mathbf{W}' \mathbf{W}] (\mathbf{W} d\mathbf{W}').$$

Proof. Taking $\boldsymbol{\mu} = 0$ and $s = (\text{tr} \mathbf{\Sigma}^{-1} \mathbf{W}' \mathbf{W})^{1/2} r$ in Theorem 3.1 we obtain the result, since

$$\begin{aligned}
& \int_0^\infty r^{m+n-1} h \left[r^2 \operatorname{tr} \boldsymbol{\Sigma}^{-1} \mathbf{W}' \mathbf{W} \right] dr \\
&= \int_0^\infty \left(\frac{s}{(\operatorname{tr} \boldsymbol{\Sigma}^{-1} \mathbf{W}' \mathbf{W})^{\frac{1}{2}}} \right) h(s^2) \frac{ds}{(\operatorname{tr} \boldsymbol{\Sigma}^{-1} \mathbf{W}' \mathbf{W})^{\frac{1}{2}}} \\
&= (\operatorname{tr} \boldsymbol{\Sigma}^{-1} \mathbf{W}' \mathbf{W})^{-\frac{m+n}{2}} \frac{\Gamma \left[\frac{m+n}{2} \right]}{2\pi^{\frac{m+n}{2}}}.
\end{aligned}$$

6 Some particular models

Finally, we give explicit shapes densities for some elliptical models.

The Kotz type I model is given by

$$h(y) = \frac{R^{T-1+\frac{K(N-1)}{2}} \Gamma \left[\frac{K(N-1)}{2} \right]}{\pi^{K(N-1)/2} \Gamma \left[T-1+\frac{K(N-1)}{2} \right]} y^{T-1} \exp\{-Ry\}.$$

Then, the corresponding k -th derivative of h , follows from $\frac{d^k}{dy^k} y^{T-1} \exp\{-Ry\}$, which is given by

$$(-R)^k y^{T-1} \exp\{-Ry\} \left\{ 1 + \sum_{m=1}^k \binom{k}{m} \left[\prod_{i=0}^{m-1} (T-1-i) \right] (-Ry)^{-m} \right\},$$

see Caro-Lopera *et al* (2009).

It is of interest the Gaussian case, i.e. when $T = 1$ and $R = \frac{1}{2}$, here the derivation is straightforward from the general density.

The required derivative follows easily, it is, $h^{(k)}(y) = \frac{R^{\frac{K(N-1)}{2}}}{\pi^{\frac{K(N-1)}{2}}} (-R)^k \exp\{-Ry\}$ and

$$\begin{aligned}
& \int_0^\infty r^{m+n+2t-1} h^{(2t)} \left[r^2 \operatorname{tr} \boldsymbol{\Sigma}^{-1} \mathbf{W}' \mathbf{W} + \operatorname{tr} \boldsymbol{\Omega} \right] (dr) \\
&= \frac{R^{\frac{M}{2}-\frac{1}{2}(m+n)+t}}{2\pi^{\frac{M}{2}}} \exp\{-R \operatorname{tr} \boldsymbol{\Omega}\} (\operatorname{tr} \boldsymbol{\Sigma}^{-1} \mathbf{W} \mathbf{W}')^{-\frac{m+n}{2}-t} \Gamma \left[\frac{m+n}{2} + t \right].
\end{aligned}$$

Hence $dF_{\mathbf{W}}(\mathbf{W})$ is given by

$$\begin{aligned}
&= \frac{\Gamma_n \left[\frac{N+K-2n}{2} \right] J(\mathbf{u}) \pi^{\frac{nK}{2}}}{2^n \pi^{\frac{n}{2}(N+K-\frac{3n-1}{2})} \Gamma_n \left[\frac{K}{2} \right] |\boldsymbol{\Sigma}|^{\frac{K}{2}}} \sum_{t=0}^\infty \sum_{\kappa} \frac{C_{\kappa}(\boldsymbol{\Omega} \boldsymbol{\Sigma}^{-1} \mathbf{W}' \mathbf{W})}{t! \left(\frac{1}{2} K \right)_{\kappa}} (\mathbf{W} d\mathbf{W}') \\
&\times \int_0^\infty r^{m+n+2t-1} h^{(2t)} \left[r^2 \operatorname{tr} \boldsymbol{\Sigma}^{-1} \mathbf{W}' \mathbf{W} + \operatorname{tr} \boldsymbol{\Omega} \right] (dr) \\
&= \frac{R^{\frac{M}{2}-\frac{1}{2}(m+n)} \Gamma_n \left[\frac{N+K-2n}{2} \right] J(\mathbf{u}) \exp\{-R \operatorname{tr} \boldsymbol{\Omega}\}}{2^{n+1} \pi^{\frac{n}{2}(N-\frac{3n-1}{2})+\frac{M}{2}} \Gamma_n \left[\frac{K}{2} \right] |\boldsymbol{\Sigma}|^{\frac{K}{2}}} \sum_{t=0}^\infty \sum_{\kappa} \frac{C_{\kappa}(R \boldsymbol{\Omega} \boldsymbol{\Sigma}^{-1} \mathbf{W}' \mathbf{W})}{t! \left(\frac{1}{2} K \right)_{\kappa}} \\
&\times (\operatorname{tr} \boldsymbol{\Sigma}^{-1} \mathbf{W} \mathbf{W}')^{-\frac{m+n}{2}-t} \Gamma \left[\frac{m+n}{2} + t \right] (\mathbf{W} d\mathbf{W}').
\end{aligned}$$

Therefore, we have proved that

Corollary 6.1. *The Gaussian SVD reflection shape density is*

$$dF_{\mathbf{W}}(\mathbf{W}) = \frac{R^{\frac{M}{2}-\frac{1}{2}(m+n)} \Gamma_n \left[\frac{N+K-2n}{2} \right] J(\mathbf{u}) \text{etr}\{-R\mathbf{\Omega}\}}{2^{n+1} \pi^{\frac{n}{2} \left(N - \frac{3n-1}{2} \right) + \frac{M}{2}} \Gamma_n \left[\frac{K}{2} \right] |\mathbf{\Sigma}|^{\frac{K}{2}}} \sum_{t=0}^{\infty} \sum_{\kappa} \frac{C_{\kappa} (R\mathbf{\Omega}\mathbf{\Sigma}^{-1}\mathbf{W}'\mathbf{W})}{t! \left(\frac{1}{2}K \right)_{\kappa}} \\ \times (\text{tr } \mathbf{\Sigma}^{-1}\mathbf{W}\mathbf{W}')^{-\frac{m+n}{2}-t} \Gamma \left[\frac{m+n}{2} + t \right] (\mathbf{W}d\mathbf{W}'), \quad (7)$$

where $M = (N-1)K$.

Finally, we propose the result for the Kotz type I model

$$h(y) = \frac{R^{T-1+\frac{K(N-1)}{2}} \Gamma \left[\frac{K(N-1)}{2} \right]}{\pi^{K(N-1)/2} \Gamma \left[T-1 + \frac{K(N-1)}{2} \right]} y^{T-1} \exp\{-Ry\},$$

Corollary 6.2. *The Kotz type I SVD reflection shape density is*

$$f_{\mathbf{W}}(\mathbf{W}) = \frac{R^{T-1+\frac{M}{2}-\frac{m+n}{2}} \Gamma \left[\frac{M}{2} \right] \Gamma_n \left[\frac{N+K-2n}{2} \right] J(\mathbf{u}) (\text{tr } \mathbf{\Omega})^{T-1} \text{etr}(-R\mathbf{\Omega})}{2^{n+1} \pi^{\frac{n}{2} \left(N - \frac{3n-1}{2} \right) + \frac{M}{2}} \Gamma_n \left[\frac{K}{2} \right] |\mathbf{\Sigma}|^{\frac{K}{2}} \Gamma \left[T-1 + \frac{M}{2} \right]} \\ \times \sum_{t=0}^{\infty} \sum_{\kappa} \frac{C_{\kappa} (R\mathbf{\Omega}\mathbf{\Sigma}^{-1}\mathbf{W}'\mathbf{W})}{t! \left(\frac{1}{2}K \right)_{\kappa}} (\mathbf{W}d\mathbf{W}') (\text{tr } \mathbf{\Sigma}^{-1}\mathbf{W}'\mathbf{W})^{-\frac{m+n}{2}-t} \\ \times \left\{ \sum_{u=0}^{\infty} \frac{\Gamma \left[\frac{M}{2} + t + u \right] \prod_{s=0}^{u-1} (T-1-s)}{u! R^u (\text{tr } \mathbf{\Omega})^u \Gamma \left[T-1 + \frac{M}{2} \right]} \right. \\ \left. + \sum_{m=1}^k \binom{k}{m} \left[\prod_{i=0}^{m-1} (T-1-i) \right] \frac{(-R)^{-m} (\text{tr } \mathbf{\Omega})^{-m}}{\Gamma \left[T-1-m + \frac{M}{2} \right]} \right. \\ \left. \times \sum_{u=0}^{\infty} \frac{\Gamma \left[\frac{M}{2} + t + u \right] \prod_{s=0}^{u-1} (T-1-m-s)}{u! R^u (\text{tr } \mathbf{\Omega})^u} \right\},$$

where $M = (N-1)K$.

Proof. As we note before the k -th derivative of h follows from,

$$\frac{d^k}{dy^k} y^{T-1} \exp\{-Ry\} \\ = (-R)^k y^{T-1} \exp\{-Ry\} \left\{ 1 + \sum_{m=1}^k \binom{k}{m} \left[\prod_{i=0}^{m-1} (T-1-i) \right] (-Ry)^{-m} \right\},$$

and the corresponding SVD reflection shape density, $dF_{\mathbf{W}}(\mathbf{W})$, is obtained after some sim-

plification as

$$\begin{aligned}
&= \frac{\Gamma_n \left[\frac{N+K-2n}{2} \right] J(\mathbf{u}) \pi^{\frac{nK}{2}}}{2^n \pi^{\frac{n}{2}(N+K-\frac{3n-1}{2})} \Gamma_n \left[\frac{K}{2} \right] |\Sigma|^{\frac{K}{2}}} \sum_{t=0}^{\infty} \sum_{\kappa} \frac{C_{\kappa} (\Omega \Sigma^{-1} \mathbf{W}' \mathbf{W})}{t! \left(\frac{1}{2} K \right)_{\kappa}} (\mathbf{W} d\mathbf{W}') \\
&\quad \times \int_0^{\infty} r^{m+n+2t-1} h^{(2t)} \left[r^2 \operatorname{tr} \Sigma^{-1} \mathbf{W}' \mathbf{W} + \operatorname{tr} \Omega \right] (dr) \\
&= \frac{R^{T-1+\frac{M}{2}-\frac{m+n}{2}} \Gamma \left[\frac{M}{2} \right] \Gamma_n \left[\frac{N+K-2n}{2} \right] J(\mathbf{u}) (\operatorname{tr} \Omega)^{T-1} \operatorname{etr}(-R\Omega)}{2^{n+1} \pi^{\frac{n}{2}(N-\frac{3n-1}{2})+\frac{M}{2}} \Gamma_n \left[\frac{K}{2} \right] |\Sigma|^{\frac{K}{2}} \Gamma \left[T-1+\frac{M}{2} \right]} \\
&\quad \times \sum_{t=0}^{\infty} \sum_{\kappa} \frac{C_{\kappa} (R\Omega \Sigma^{-1} \mathbf{W}' \mathbf{W})}{t! \left(\frac{1}{2} K \right)_{\kappa}} (\mathbf{W} d\mathbf{W}') (\operatorname{tr} \Sigma^{-1} \mathbf{W}' \mathbf{W})^{-\frac{m+n}{2}-t} \\
&\quad \times \left\{ \sum_{u=0}^{\infty} \frac{\Gamma \left[\frac{M}{2} + t + u \right] \prod_{s=0}^{u-1} (T-1-s)}{u! R^u (\operatorname{tr} \Omega)^u \Gamma \left[T-1+\frac{M}{2} \right]} \right. \\
&\quad + \sum_{m=1}^k \binom{k}{m} \left[\prod_{i=0}^{m-1} (T-1-i) \right] \frac{(-R)^{-m} (\operatorname{tr} \Omega)^{-m}}{\Gamma \left[T-1-m+\frac{M}{2} \right]} \\
&\quad \left. \times \sum_{u=0}^{\infty} \frac{\Gamma \left[\frac{M}{2} + t + u \right] \prod_{s=0}^{u-1} (T-1-m-s)}{u! R^u (\operatorname{tr} \Omega)^u} \right\}. \quad \square
\end{aligned}$$

The Gaussian case can be derived again by taking $T = 1$ in the above result.

6.1 Example: Mouse Vertebra

This classical application is studied in the Gaussian case by Dryden and Mardia (1998). Here we consider again the same model and contrasted it, via the modified BIC criterion, with two non Gaussian models.

The isotropic Gaussian shape density is given by

$$\begin{aligned}
dF_{\mathbf{W}}(\mathbf{W}) &= \frac{2^{-\frac{1}{2}(2-m+M+n)} \Gamma_n \left[\frac{N+K-2n}{2} \right] J(\mathbf{u})}{\pi^{\frac{1}{4}(2M+n-3n^2+2nM)} \sigma^{-(m-M+n)} \Gamma_n \left[\frac{K}{2} \right]} \operatorname{etr} \left(-\frac{\boldsymbol{\mu}' \boldsymbol{\mu}}{2\sigma^2} \right) \\
&\quad \times \sum_{t=0}^{\infty} \frac{\Gamma \left[\frac{m+n}{2} + t \right]}{t!} \sum_{\kappa} \frac{C_{\kappa} \left(\frac{1}{2\sigma^2} \boldsymbol{\mu}' \mathbf{W} \mathbf{W}' \boldsymbol{\mu} \right)}{\left(\frac{1}{2} K \right)_{\kappa}} (\mathbf{W} d\mathbf{W}'), \quad (8)
\end{aligned}$$

where $M = K(N-1)$, $n = \min\{(N-1), K\}$ and $m = (N-1)n - 1$. Here we study three models, the Gaussian shape (N), and the Kotz (K) model for $T = 2$ and $T = 3$.

The shape density associated to the Kotz model indexed by $T = 2$, $R = \frac{1}{2}$ (and $s = 1$) is given by:

$$\begin{aligned}
dF_{\mathbf{W}}(\mathbf{W}) &= \frac{2^{-\frac{1}{2}(-m+M+n)} \Gamma_n \left[\frac{N+K-2n}{2} \right] J(\mathbf{u})}{\pi^{\frac{1}{4}(2M+n-3n^2+2nM)} \sigma^{-(m-M+n)} M \Gamma_n \left[\frac{K}{2} \right]} \text{etr} \left(-\frac{\boldsymbol{\mu}' \boldsymbol{\mu}}{2\sigma^2} \right) \\
&\times \sum_{t=0}^{\infty} \frac{\left(\text{tr} \left(\frac{\boldsymbol{\mu}' \boldsymbol{\mu}}{2\sigma^2} \right) - 2t \right) \Gamma \left[\frac{m+n}{2} + t \right] + \Gamma \left[\frac{m+n}{2} + t + 1 \right]}{t!} \\
&\times \sum_{\kappa} \frac{C_{\kappa} \left(\frac{1}{2\sigma^2} \boldsymbol{\mu}' \mathbf{W} \mathbf{W}' \boldsymbol{\mu} \right)}{\left(\frac{1}{2} K \right)_{\kappa}} (\mathbf{W} d\mathbf{W}').
\end{aligned} \tag{9}$$

And the corresponding density, $dF_{\mathbf{W}}(\mathbf{W})$, for the Kotz model $T = 3$, is obtained as:

$$\begin{aligned}
&= \frac{2^{-\frac{1}{2}(-2-m+M+n)} \Gamma_n \left[\frac{N+K-2n}{2} \right] J(\mathbf{u})}{\pi^{\frac{1}{4}(2M+n-3n^2+2nM)} \sigma^{-(m-M+n)} M(M+2) \Gamma_n \left[\frac{K}{2} \right]} \text{etr} \left(-\frac{\boldsymbol{\mu}' \boldsymbol{\mu}}{2\sigma^2} \right) \\
&\times \sum_{t=0}^{\infty} \left\{ \left[4t^2 - 2t - 4t \text{tr} \left(\frac{\boldsymbol{\mu}' \boldsymbol{\mu}}{2\sigma^2} \right) + \text{tr}^2 \left(\frac{\boldsymbol{\mu}' \boldsymbol{\mu}}{2\sigma^2} \right) \right] \Gamma \left[\frac{m+n}{2} + t \right] \right. \\
&\quad \left. + \left[-4t + 2 \text{tr} \left(\frac{\boldsymbol{\mu}' \boldsymbol{\mu}}{2\sigma^2} \right) \right] \Gamma \left[\frac{m+n}{2} + t + 1 \right] + \Gamma \left[\frac{m+n}{2} + t + 2 \right] \right\} \\
&\times \sum_{\kappa} \frac{C_{\kappa} \left(\frac{1}{2\sigma^2} \boldsymbol{\mu}' \mathbf{W} \mathbf{W}' \boldsymbol{\mu} \right)}{t! \left(\frac{1}{2} K \right)_{\kappa}} (\mathbf{W} d\mathbf{W}').
\end{aligned}$$

In order to decide which the elliptical model is the best one, different criteria have been employed for the model selection. We shall consider a modification of the BIC statistic as discussed in Yang and Yang (2007), and which was first achieved by Rissanen (1978) in a coding theory framework. The modified BIC is given by:

$$BIC^* = -2\mathfrak{L}(\tilde{\boldsymbol{\mu}}, \tilde{\sigma}^2, h) + n_p(\log(n+2) - \log 24),$$

where $\mathfrak{L}(\tilde{\boldsymbol{\mu}}, \tilde{\sigma}^2, h)$ is the maximum of the log-likelihood function, n is the sample size and n_p is the number of parameters to be estimated for each particular shape density.

As proposed by Kass and Raftery (1995) and Raftery (1995), the following selection criteria have been employed for the model selection.

Table 1: Grades of evidence corresponding to values of the BIC^* difference.

BIC^* difference	Evidence
0–2	Weak
2–6	Positive
6–10	Strong
> 10	Very strong

Fixing the variance of the process as 50 (the maximum median variances of the two samples), the maximum likelihood estimators for location parameters associated with the small and large groups are summarized in the following table:

According to the modified BIC criterion, the Kotz model with parameters $T = 3$, $R = \frac{1}{2}$ and $s = 1$ is the most appropriate among the three elliptical densities for modeling the data. There is a very strong difference between the non Gaussian and the classical Gaussian model in this experiment.

Table 2: The maximum likelihood estimators

Group	BIC^* $K : T = 3$ $K : T = 2$	$\tilde{\mu}_{11}$	$\tilde{\mu}_{12}$	$\tilde{\mu}_{21}$	$\tilde{\mu}_{22}$	$\tilde{\mu}_{31}$
Small	$\begin{matrix} -5.9146 \\ -39.6272 \\ -23.0250 \end{matrix}$	$\begin{matrix} -1.9214 \\ 2.1250 \\ -6.1682 \end{matrix}$	$\begin{matrix} -42.5338 \\ -47.8016 \\ -45.0331 \end{matrix}$	$\begin{matrix} 14.1761 \\ 16.3513 \\ 14.6035 \end{matrix}$	$\begin{matrix} -4.8190 \\ -3.9691 \\ -6.4983 \end{matrix}$	$\begin{matrix} 24.0766 \\ 26.9744 \\ 25.5710 \end{matrix}$
Large	$\begin{matrix} 24.5000 \\ -9.2156 \\ 7.3880 \end{matrix}$	$\begin{matrix} -23.1834 \\ -34.4190 \\ -42.9562 \end{matrix}$	$\begin{matrix} -32.9246 \\ -29.3792 \\ -1.6455 \end{matrix}$	$\begin{matrix} 8.1455 \\ 5.9227 \\ -3.6885 \end{matrix}$	$\begin{matrix} -10.5612 \\ -13.7687 \\ -13.7522 \end{matrix}$	$\begin{matrix} 22.5117 \\ 20.9197 \\ 3.6578 \end{matrix}$

$\tilde{\mu}_{32}$	$\tilde{\mu}_{41}$	$\tilde{\mu}_{42}$	$\tilde{\mu}_{51}$	$\tilde{\mu}_{52}$
$\begin{matrix} -0.2605 \\ 2.1288 \\ -2.6121 \end{matrix}$	$\begin{matrix} 4.8195 \\ 4.9123 \\ 5.5892 \end{matrix}$	$\begin{matrix} 4.7981 \\ 5.8552 \\ 4.6337 \end{matrix}$	$\begin{matrix} -29.2691 \\ -33.0646 \\ -30.8241 \end{matrix}$	$\begin{matrix} -0.7072 \\ 0.4382 \\ 6.0498 \end{matrix}$
$\begin{matrix} -12.8674 \\ -20.2846 \\ -27.4382 \end{matrix}$	$\begin{matrix} 5.0300 \\ 6.1263 \\ 5.2368 \end{matrix}$	$\begin{matrix} 2.3324 \\ 1.1366 \\ -2.7585 \end{matrix}$	$\begin{matrix} -26.0251 \\ -23.1221 \\ -0.9942 \end{matrix}$	$\begin{matrix} 18.6865 \\ 27.5996 \\ 34.1886 \end{matrix}$

Let μ_1 and μ_2 be the mean shape of the small and large groups, respectively. We test equal mean shape under the best model, and the likelihood ratio (based on $-2 \log \Lambda \approx \chi^2_{10}$) for the test $H_0 : \mu_1 = \mu_2$ vs $H_a : \mu_1 \neq \mu_2$, provides the p-value 0.84, which means that there are extremely evidence that the mean shapes of the two groups are equal if the variance of the experiment is fixed in 50 (the maximum median of the variances of the two samples), a deeper study of this case is suggested, because the variance estimation was problematic in the performed inference procedure for these data. We highlight that our intention is to illustrate the technique and performed inference with an exact likelihood efficiently computable after modification of the algorithms given for hypergeometric series (Koev and Edelman (2006)).

A final comment, for any elliptical model we can obtain the SVD reflection model, however a nontrivial problem appears, the $2t$ -th derivative of the generator model, which can be seen as a partition theory problem. For The general case of a Kotz model ($s \neq 1$), and another models like Pearson II and VII, Bessel, Jensen-logistic, we can use formulae for these derivatives given by Caro-Lopera *et al* (2009). The resulting densities have again a form of a generalized series of zonal polynomials which can be computed efficiently after some modification of existing works for hypergeometric series (see Koev and Edelman (2006)), thus the inference over an exact density can be performed, avoiding the use of any asymptotic distribution, and the initial transformation avoids the invariant polynomials of Davis (1908), and it lets the inclusion of any correlation among landmarks.

Acknowledgment

This research work was supported by University of Medellin (Medellin, Colombia) and Universidad Autónoma Agraria Antonio Narro (México), joint grant No. 469, SUMMA group. Also, the first author was partially supported by CONACYT - México, research grant no. 138713 and IDI-Spain, Grants No. FQM2006-2271 and MTM2008-05785 and the paper was written during J. A. Díaz- García's stay as a visiting professor at the Department of Statistics and O. R. of the University of Granada, Spain.

References

F. J. Caro-Lopera, J. A. Díaz-García and G. González-Farías (2009). Noncentral elliptical configuration density. J. Multivariate Anal. 101(1), 32–43.

- A. W. Davis (1980). Invariant polynomials with two matrix arguments, extending the zonal polynomials. In: *Multivariate Analysis V*, (Krishnaiah, P. R. ed.), North-Holland.
- J. A. Díaz-García, R. Gutiérrez- Jáimez, and K. V. Mardia (1997). Wishart and Pseudo-Wishart distributions and some applications to shape theory. *J. Multivariate Anal.* 63, 73-87.
- J. A. Díaz-García, J. R. Gutiérrez, and R. Ramos (2003). Size-and-Shape Cone, Shape Disk and Configuration Densities for the Elliptical Models. *Braz. J. Probab. Stat.* 17, 135-146.
- I. L. Dryden and K.V. Mardia, *Statistical shape analysis*. John Wiley and Sons, Chichester, 1998.
- K. T. Fang, and Y. T. Zhang, *Generalized Multivariate Analysis*. Science Press, Springer-Verlag, Beijing, 1990.
- C. R. Goodall (1991). Procrustes methods in the statistical analysis of shape (with discussion). *J. Roy. Statist. Soc. Ser. B.* 53, 285-339.
- C. R. Goodall, and K. V. Mardia (1991). A geometrical derivation of the shape density. *Adv. in Appl. Probab.* 23, 496-514.
- C. R. Goodall, and K. V. Mardia (1993). Multivariate Aspects of Shape Theory. *Ann. Statist.* 21, 848-866.
- A. K. Gupta, and T. Varga, *Elliptically Contoured Models in Statistics*. Kluwer Academic Publishers, Dordrecht, 1993.
- A. T. James (1964). Distributions of matrix variate and latent roots derived from normal samples. *Ann. Math. Statist.* 35, 475-501.
- R. E. Kass, and A. E. Raftery (1995). Bayes factor. *J. Amer. Statist. Soc.* 90, 773-795.
- P. Koev and A. Edelman (2006). The efficient evaluation of the hypergeometric function of a matrix argument. *Math. Comp.* 75, 833-846.
- H. L. Le and D. G. Kendall (1993). The Riemannian structure of Euclidean spaces: a novel environment for statistics. *Ann. Statist.* 21, 1225-1271.
- R. J. Muirhead, *Aspects of multivariate statistical theory*. Wiley Series in Probability and Mathematical Statistics. John Wiley & Sons, Inc. 1982.
- A. E. Raftery (1995). Bayesian model selection in social research. *Sociological Methodology*, 25, 111-163.
- J. Rissanen (1978). Modelling by shortest data description. *Automatica*, 14, 465-471.
- M. S. Srivastava, and C. G. Khatri, *An Introduction to Multivariate Statistics*. North Holland, 1979.
- Ch. Ch. Yang, and Ch. Ch. Yang (2007). Separating latent classes by information criteria. *J. Classification* 24, 183-203.